Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.

Antibiotics For Patients With Total Joint Replacements Who Are Having Dental And Other Procedures

Bacterial infections should be treated promptly to avoid seeding of the artificial joint. This patient should receive antibiotic prophylaxis prior to any procedure which is likely to cause bacteremia. Guidelines regarding prophylactics can be found in **JAMA**, December 12, 1990, Vol 264, No. 22, Pages 2919 - 2922. Also, referenced in the **AAOS Bulletin**, July 1997.

For dental procedures this should be **3.0 grams AMOXICILLIN orally 1 hour before the procedure, and 1.5 grams 6 hours later**. If Allergic to PCN, use **CLINDAMYCIN 600 mg P.O. 1 hour before procedure and 150 mg P.O. 6 hours later.** Use appropriate antibiotics for other procedures.